

Stay Current on RSV Immunization Guidance

Pharmacies are preparing for another cold and flu season involving respiratory syncytial virus (RSV).

RSV is the most common cause of infant hospitalizations in the US. Older adults also have a high risk of complications (intubation, etc).

Unvaccinated older adults are eligible for immunization if they are 75 and up...or 50 to 74 with certain risk factors (COPD, diabetes, etc).

Recommend any of the adult vaccines...Abrysvo, Arexvy, or mResvia. Compare options with our *RSV Vaccines* chart. Ideally vaccinate in late summer or early fall depending on when your region's RSV season starts.

Be aware that FDA approved Abrysvo and mResvia for high-risk adults between 18 and 49 years. But the Advisory Committee on Immunization Practices doesn't recommend routine RSV vaccination for these patients until we have more safety and efficacy data.

Advise Abrysvo for unvaccinated moms between 32 and 36 weeks of pregnancy...in September to January...to pass antibodies to their baby.

But avoid Arexvy and mResvia...they are NOT approved in pregnancy.

For moms vaccinated this season, share that their newborn won't need their own RSV immunization after delivery.

On the other hand, their baby should get clesrovimab (Enflonsia) or nirsevimab (Beyfortus)...ideally within a week after birth...if mom delivers within 2 weeks after getting Abrysvo or if the dose was in a PRIOR pregnancy. Get our *RSV Monoclonal Antibodies* chart for dosing, etc.

Compare RSV product differences to avoid errors.

For example, point out that Abrysvo and Arexvy are refrigerated while mResvia is frozen. And only Abrysvo and Arexvy are reconstituted.

Watch for Abrysvo product changes. The vial and diluent syringe kits are being phased out. This season, expect to only get the dual-chamber Act-O-Vial version that mixes drug and diluent when activated.

Limit confusion by ideally stocking the fewest vaccine options needed to cover your patient population. For example, consider prioritizing Abrysvo if you serve both pregnant and older adults.

Clarify that we don't have evidence or recommendations yet for giving adults more than 1 vaccine dose. Check vaccine records carefully to ensure adults only get 1 lifetime dose for now.

Use our *Preventing RSV* chart to answer other common questions about RSV risk factors, who should get a vaccine, and more.

Key References:

- CDC. RSV Vaccine Guidance for Adults. July 8, 2025. <https://www.cdc.gov/rsv/hcp/vaccine-clinical-guidance/adults.html> (Accessed September 16, 2025).
- The American College of Obstetricians and Gynecologists. Maternal Respiratory Syncytial Virus Vaccination. August 2025. <https://www.acog.org/clinical/clinical-guidance/practice-advisory/articles/2023/09/maternal-respiratory-syncytial-virus-vaccination> (Accessed September 16, 2025).
- Committee on Infectious Diseases. Recommendations for the Prevention of RSV Disease in Infants and Children: Policy Statement. *Pediatrics*. 2025 Aug 19. doi: 10.1542/peds.2025-073923.

Pharmacist's Letter. October 2025, No. 411004

Cite this document as follows: Article, Stay Current on RSV Immunization Guidance, Hospital Pharmacy Technician's Letter, October 2025

The content of this article is provided for educational and informational purposes only, and is not a substitute for the advice, opinion or diagnosis of a trained medical professional. If your organization is interested in an enterprise subscription, email sales@trchealthcare.com.

© 2025 Therapeutic Research Center (TRC). TRC and Hospital Pharmacy Technician's Letter and the associated logo(s) are trademarks of Therapeutic Research Center. All Rights Reserved.